

November 17, 2016

6.453 *Quantum Optical Communication* Lecture 19

Jeffrey H. Shapiro

Optical and Quantum Communications Group

RESEARCH LABORATORY OF ELECTRONICS
Massachusetts Institute of Technology

www.rle.mit.edu/qoptics

6.453 *Quantum Optical Communication* - Lecture 19

- Announcements
 - Pick up lecture notes, slides
- Continuous-Time Photodetection
 - Noise spectral densities in direct detection
 - Semiclassical theory of coherent detection
 - Quantum theory of coherent detection
 - Coherent-detection signatures of non-classical light

Semiclassical versus Quantum Photodetection

- Semiclassical Theory: Given $\{P(t) : -\infty < t < \infty\}$
 - $\{i(t) : -\infty < t < \infty\}$ is an inhomogeneous Poisson Impulse Train
 - Rate function $\lambda(t) \equiv \eta P(t)/\hbar\omega_o$

- Quantum Theory:

$$\hat{E}'(t) \equiv \sqrt{\eta} \hat{E}(t) + \sqrt{1-\eta} \hat{E}_\eta(t)$$

$$i(t) \leftrightarrow \hat{i}(t) \equiv q \hat{E}'^\dagger(t) \hat{E}'(t)$$

Stationary Statistics for Continuous-Wave Sources

- Stationary Mean and Covariance Functions:

$$\langle x(t) \rangle = \text{constant} \equiv \langle x \rangle$$

$$\langle \Delta x(t+\tau) \Delta x(t) \rangle = \text{function of } \tau \text{ only} \equiv K_{xx}(\tau)$$

- Semiclassical Photodetection:

$$\langle i \rangle = q \frac{\eta \langle P \rangle}{\hbar\omega_o} \quad \text{and} \quad K_{ii}(\tau) = q \langle i \rangle \delta(\tau) + q^2 \frac{\eta^2 K_{PP}(\tau)}{(\hbar\omega_o)^2}$$

- Quantum Photodetection:

$$\langle i \rangle = q\eta \langle \hat{E}^\dagger(0) \hat{E}(0) \rangle$$


$$K_{ii}(\tau) = q \langle i \rangle \delta(\tau) + q^2 \eta^2 [\langle \hat{E}^\dagger(\tau) \hat{E}^\dagger(0) \hat{E}(\tau) \hat{E}(0) \rangle - \langle \hat{E}^\dagger(0) \hat{E}(0) \rangle^2]$$

Photocurrent Noise Spectral Density

- Photocurrent Noise Spectral Density:

$$\mathcal{S}_{ii}(\omega) \equiv \int_{-\infty}^{\infty} d\tau K_{ii}(\tau) e^{-j\omega\tau}$$

- Propagation Through a Linear Time-Invariant Filter:



$$\mathcal{S}_{i'i'}(\omega) = \mathcal{S}_{ii}(\omega) |H(\omega)|^2$$

- Physical Interpretation: $\mathcal{S}_{ii}(\omega)$ = mean-squared fluctuation strength per unit bilateral bandwidth (in Hz) in frequency ω components of $i(t)$

Direct-Detection Signatures of Non-Classical Light

- Semiclassical Theory:

$$\mathcal{S}_{ii}(\omega) = q\langle i \rangle + q^2 \frac{\eta^2 \mathcal{S}_{PP}(\omega)}{(\hbar\omega_o)^2} \geq q\langle i \rangle$$


- Quantum Theory:

$$\mathcal{S}_{ii}(\omega) \geq 0$$

- Sub-Shot-Noise Non-Classical Signature:

$$\mathcal{S}_{ii}(\omega) < q\langle i \rangle$$

Balanced Homodyne Detection



- Signal = $E(t)$ or $\hat{E}(t)$, LO = $\sqrt{\frac{P_{LO}}{\hbar\omega_o}} e^{j\theta}$ or $\left| \sqrt{\frac{P_{LO}}{\hbar\omega_o}} e^{j\theta} \right\rangle$
- Low-Pass Filter: $H(\omega) = 1$, for $|\omega| \leq \frac{\Delta\omega}{2}$

Balanced Homodyne Detection (Within Passband)

- Semiclassical Statistics in Strong Local Oscillator Limit:

$$i_{\text{hom}}(t) = 2q\eta\sqrt{\frac{P_{LO}}{\hbar\omega_o}} \text{Re}(E(t)e^{-j\theta}) + i_{LO}(t)$$

- Gaussian-process local oscillator shot noise:

$$\langle i_{LO} \rangle = 0 \quad \text{and} \quad \mathcal{S}_{i_{LO}i_{LO}}(\omega) = q^2 \frac{\eta P_{LO}}{\hbar\omega_o}$$


- Quantum Statistics in Strong Local Oscillator Limit:

$$i_{\text{hom}}(t) \leftrightarrow \hat{i}_{\text{hom}}(t) = 2q\eta\sqrt{\frac{P_{LO}}{\hbar\omega_o}} \text{Re}(\hat{E}(t)e^{-j\theta}) + i_{\eta}(t)$$

- Gaussian-process sub-unity quantum efficiency noise:

$$\langle i_{\eta} \rangle = 0 \quad \text{and} \quad \mathcal{S}_{i_{\eta}i_{\eta}}(\omega) = q^2(1 - \eta) \frac{\eta P_{LO}}{\hbar\omega_o}$$

Balanced Heterodyne Detection



- Signal = $E_S(t)e^{-j\omega_{IF}t}$ or $\hat{E}_S(t)e^{-j\omega_{IF}t} + \hat{E}_I(t)e^{j\omega_{IF}t}$
- LO = $\sqrt{P_{LO}/\hbar\omega_o}$ or $|\sqrt{P_{LO}/\hbar\omega_o}\rangle$
- Bandpass Filter: $H(\omega) = 1$, for $|\omega \pm \omega_{IF}| \leq \frac{\Delta\omega}{2}$

Balanced Heterodyne Detection (Within Passband)

- Semiclassical Statistics in Strong Local Oscillator Limit:

$$i_{\text{het}}(t) = 2q\eta\sqrt{\frac{P_{LO}}{\hbar\omega_o}}\text{Re}(E_S(t)e^{-j\omega_{IF}t}) + i_{LO}(t)$$

- Gaussian-process local oscillator shot noise:

$$\langle i_{LO} \rangle = 0 \quad \text{and} \quad \mathcal{S}_{i_{LO}i_{LO}}(\omega) = q^2\frac{\eta P_{LO}}{\hbar\omega_o}$$

- Quantum Statistics in Strong Local Oscillator Limit:

$$i_{\text{het}}(t) \leftrightarrow \hat{i}_{\text{het}}(t) = 2q\eta\sqrt{\frac{P_{LO}}{\hbar\omega_o}}\text{Re}[(\hat{E}_S(t) + \hat{E}_I^\dagger(t))e^{-j\omega_{IF}t}] + i_\eta(t)$$

- Gaussian-process sub-unity quantum efficiency noise:

$$\langle i_\eta \rangle = 0 \quad \text{and} \quad \mathcal{S}_{i_\eta i_\eta}(\omega) = q^2(1 - \eta)\frac{\eta P_{LO}}{\hbar\omega_o}$$

Coherent-Detection Non-Classical Light Signatures

- Semiclassical Theory (within filter passband):

$$\mathcal{S}_{i_{\text{hom}}i_{\text{hom}}}(\omega) \geq \mathcal{S}_{i_{\text{LO}}i_{\text{LO}}}(\omega) = q^2 \frac{\eta P_{\text{LO}}}{\hbar\omega_o}$$

$$\mathcal{S}_{i_{\text{het}}i_{\text{het}}}(\omega) \geq \mathcal{S}_{i_{\text{LO}}i_{\text{LO}}}(\omega) = q^2 \frac{\eta P_{\text{LO}}}{\hbar\omega_o}$$

- Quantum Theory (within filter passband):

$$\mathcal{S}_{i_{\text{hom}}i_{\text{hom}}}(\omega) \geq \mathcal{S}_{i_{\eta}i_{\eta}}(\omega) = \frac{q^2 \eta (1 - \eta) P_{\text{LO}}}{\hbar\omega_o}$$

- Sub-Shot-Noise Spectra (in passband) are Non-Classical:

$$\mathcal{S}_{i_{\text{hom}}i_{\text{hom}}}(\omega) \text{ or } \mathcal{S}_{i_{\text{het}}i_{\text{het}}}(\omega) < q^2 \frac{\eta P_{\text{LO}}}{\hbar\omega_o}$$

Coming Attractions: Lecture 20

- Lecture 20:

Nonlinear Optics of $\chi^{(2)}$ Interactions

- Maxwell's equations with a nonlinear polarization
- Coupled-mode equations for parametric downconversion
- Phase-matching for efficient interactions

MIT OpenCourseWare
<https://ocw.mit.edu>

6.453 Quantum Optical Communication
Fall 2016

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.