

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.102 Introduction to Functional Analysis  
Spring 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

SOLUTIONS TO PROBLEM SET 10

*Problem P10.1* Let  $H$  be a separable, infinite dimensional Hilbert space. Show that the direct sum of two copies of  $H$  is a Hilbert space with the norm

$$(23.18) \quad H \oplus H \ni (u_1, u_2) \mapsto (\|u_1\|_H^2 + \|u_2\|_H^2)^{\frac{1}{2}}$$

either by constructing an isometric isomorphism

$$(23.19) \quad T : H \longrightarrow H \oplus H, \text{ 1-1 and onto, } \|u\|_H = \|Tu\|_{H \oplus H}$$

or otherwise. In any case, construct a map as in (23.19).

Solution: Let  $\{e_i\}_{i \in \mathbb{N}}$  be an orthonormal basis of  $H$ , which exists by virtue of the fact that it is an infinite-dimensional but separable Hilbert space. Define the map

$$(23.20) \quad T : H \ni u \longrightarrow \left( \sum_{i=1}^{\infty} (u, e_{2i-1}) e_i, \sum_{i=1}^{\infty} (u, e_{2i}) e_i \right) \in H \oplus H$$

The convergence of the Fourier Bessel series shows that this map is well-defined and linear. Injectivity similarly follows from the fact that  $Tu = 0$  in the image implies that  $(u, e_i) = 0$  for all  $i$  and hence  $u = 0$ . Surjectivity is also clear from the fact that

$$(23.21) \quad S : H \oplus H \ni (u_1, u_2) \mapsto \sum_{i=1}^{\infty} ((u_1, e_i) e_{2i-1} + (u_2, e_i) e_{2i}) \in H$$

is a 2-sided inverse and Bessel's identity implies isometry since  $\|S(u_1, u_2)\|^2 = \|u_1\|^2 + \|u_2\|^2$

*Problem P10.2* One can repeat the preceding construction any finite number of times. Show that it can be done 'countably often' in the sense that if  $H$  is a separable, infinite dimensional, Hilbert space then

$$(23.22) \quad l_2(H) = \left\{ u : \mathbb{N} \longrightarrow H; \|u\|_{l_2(H)}^2 = \sum_i \|u_i\|_H^2 < \infty \right\}$$

has a Hilbert space structure and construct an explicit isometric isomorphism from  $l_2(H)$  to  $H$ .

Solution: A similar argument as in the previous problem works. Take an orthonormal basis  $e_i$  for  $H$ . Then the elements  $E_{i,j} \in l_2(H)$ , which for each  $i, j$  consist of the sequences with 0 entries except the  $j$ th, which is  $e_i$ , given an orthonormal basis for  $l_2(H)$ . Orthogonality is clear, since with the inner product is

$$(23.23) \quad (u, v)_{l_2(H)} = \sum_j (u_j, v_j)_H.$$

Completeness follows from completeness of the orthonormal basis of  $H$  since if  $v = \{v_j\}$   $(v, E_{j,i}) = 0$  for all  $j$  implies  $v_j = 0$  in  $H$ . Now, to construct an isometric isomorphism just choose an isomorphism  $m : \mathbb{N}^2 \longrightarrow \mathbb{N}$  then

$$(23.24) \quad Tu = v, \quad v_j = \sum_i (u, e_{m(i,j)}) e_i \in H.$$

I would expect you to go through the argument to check injectivity, surjectivity and that the map is isometric.

*Problem P10.3* Recall, or perhaps learn about, the winding number of a closed curve with values in  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ . We take as given the following fact:<sup>4</sup> If  $Q = [0, 1]^N$  and  $f : Q \rightarrow \mathbb{C}^*$  is continuous then for each choice of  $b \in \mathbb{C}$  satisfying  $\exp(2\pi i b) = f(0)$ , there exists a unique continuous function  $F : Q \rightarrow \mathbb{C}$  satisfying

$$(23.25) \quad \exp(2\pi i F(q)) = f(q), \quad \forall q \in Q \text{ and } F(0) = b.$$

Of course, you are free to change  $b$  to  $b + n$  for any  $n \in \mathbb{Z}$  but then  $F$  changes to  $F + n$ , just shifting by the same integer.

- (1) Now, suppose  $c : [0, 1] \rightarrow \mathbb{C}^*$  is a closed curve – meaning it is continuous and  $c(1) = c(0)$ . Let  $C : [0, 1] \rightarrow \mathbb{C}$  be a choice of  $F$  for  $N = 1$  and  $f = c$ . Show that the winding number of the closed curve  $c$  may be defined *unambiguously* as

$$(23.26) \quad \text{wn}(c) = C(1) - C(0) \in \mathbb{Z}.$$

Solution: Let  $C'$ , be another choice of  $F$  in this case. Now,  $g(t) = C'(t) - C(t)$  is continuous and satisfies  $\exp(2\pi i g(t)) = 1$  for all  $t \in [0, 1]$  so by the uniqueness must be constant, thus  $C'(1) - C'(0) = C(1) - C(0)$  and the winding number is well-defined.

- (2) Show that  $\text{wn}(c)$  is *constant under homotopy*. That is if  $c_i : [0, 1] \rightarrow \mathbb{C}^*$ ,  $i = 1, 2$ , are two closed curves so  $c_i(1) = c_i(0)$ ,  $i = 1, 2$ , which are *homotopic* through closed curves in the sense that there exists  $f : [0, 1]^2 \rightarrow \mathbb{C}^*$  continuous and such that  $f(0, x) = c_1(x)$ ,  $f(1, x) = c_2(x)$  for all  $x \in [0, 1]$  and  $f(y, 0) = f(y, 1)$  for all  $y \in [0, 1]$ , then  $\text{wn}(c_1) = \text{wn}(c_2)$ .

Solution: Choose  $F$  using the ‘fact’ corresponding to this homotopy  $f$ . Since  $f$  is periodic in the second variable – the two curves  $f(y, 0)$ , and  $f(y, 1)$  are the same – so by the uniqueness  $F(y, 0) - F(y, 1)$  must be constant, hence  $\text{wn}(c_2) = F(1, 1) - F(1, 0) = F(0, 1) - F(0, 0) = \text{wn}(c_1)$ .

- (3) Consider the closed curve  $L_n : [0, 1] \ni x \mapsto e^{2\pi i x} \text{Id}_{n \times n}$  of  $n \times n$  matrices. Using the standard properties of the determinant, show that this curve is not homotopic to the identity through closed curves in the sense that there does not exist a continuous map  $G : [0, 1]^2 \rightarrow \text{GL}(n)$ , with values in the invertible  $n \times n$  matrices, such that  $G(0, x) = L_n(x)$ ,  $G(1, x) \equiv \text{Id}_{n \times n}$  for all  $x \in [0, 1]$ ,  $G(y, 0) = G(y, 1)$  for all  $y \in [0, 1]$ .

Solution: The determinant is a continuous (actually it is analytic) map which vanishes precisely on non-invertible matrices. Moreover, it is given by the product of the eigenvalues so

$$(23.27) \quad \det(L_n) = \exp(2\pi i x n).$$

This is a periodic curve with winding number  $n$  since it has the ‘lift’  $xn$ . Now, if there were to exist such an homotopy of periodic curves of matrices, always invertible, then by the previous result the winding number of the determinant would have to remain constant. Since the winding number for the constant curve with value the identity is 0 such an homotopy cannot exist.

*Problem P10.4* Consider the closed curve corresponding to  $L_n$  above in the case of a separable but now infinite dimensional Hilbert space:

$$(23.28) \quad L : [0, 1] \ni x \mapsto e^{2\pi i x} \text{Id}_H \in \text{GL}(H) \subset \mathcal{B}(H)$$

---

<sup>4</sup>Of course, you are free to give a proof – it is not hard.

taking values in the invertible operators on  $H$ . Show that after identifying  $H$  with  $H \oplus H$  as above, there is a continuous map

$$(23.29) \quad M : [0, 1]^2 \longrightarrow \text{GL}(H \oplus H)$$

with values in the invertible operators and satisfying

$$(23.30) \quad M(0, x) = L(x), \quad M(1, x)(u_1, u_2) = (e^{4\pi i x} u_1, u_2), \quad M(y, 0) = M(y, 1), \quad \forall x, y \in [0, 1].$$

Hint: So, think of  $H \oplus H$  as being 2-vectors  $(u_1, u_2)$  with entries in  $H$ . This allows one to think of ‘rotation’ between the two factors. Indeed, show that

$$(23.31) \quad U(y)(u_1, u_2) = (\cos(\pi y/2)u_1 + \sin(\pi y/2)u_2, -\sin(\pi y/2)u_1 + \cos(\pi y/2)u_2)$$

defines a continuous map  $[0, 1] \ni y \mapsto U(y) \in \text{GL}(H \oplus H)$  such that  $U(0) = \text{Id}$ ,  $U(1)(u_1, u_2) = (u_2, -u_1)$ . Now, consider the 2-parameter family of maps

$$(23.32) \quad U^{-1}(y)V_2(x)U(y)V_1(x)$$

where  $V_1(x)$  and  $V_2(x)$  are defined on  $H \oplus H$  as multiplication by  $\exp(2\pi i x)$  on the first and the second component respectively, leaving the other fixed.

Solution: Certainly  $U(y)$  is invertible since its inverse is  $U(-y)$  as follows in the two dimensional case. Thus the map  $W(x, y)$  on  $[0, 1]^2$  in (23.32) consists of invertible and bounded operators on  $H \oplus H$ , meaning a continuous map  $W : [0, 1]^2 \longrightarrow \text{GL}(H \oplus H)$ . When  $x = 0$  or  $x = 1$ , both  $V_1(x)$  and  $v_2(x)$  reduce to the identity, and hence  $W(0, y) = W(1, y)$  for all  $y$ , so  $W$  is periodic in  $x$ . Moreover at  $y = 0$   $W(x, 0) = V_2(x)V_1(x)$  is exactly  $L(x)$ , a multiple of the identity. On the other hand, at  $x = 1$  we can track composite as

$$(23.33) \quad \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \mapsto \begin{pmatrix} e^{2\pi i x} u_1 \\ u_2 \end{pmatrix} \mapsto \begin{pmatrix} u_2 \\ -e^{2\pi i x} u_1 \end{pmatrix} \mapsto \begin{pmatrix} u_2 \\ -e^{4\pi i x} u_1 \end{pmatrix} \mapsto \begin{pmatrix} e^{4\pi i x} u_1 \\ u_2 \end{pmatrix}.$$

This is what is required of  $M$  in (23.30).

*Problem P10.5* Using a rotation similar to the one in the preceding problem (or otherwise) show that there is a continuous map

$$(23.34) \quad G : [0, 1]^2 \longrightarrow \text{GL}(H \oplus H)$$

such that

$$(23.35) \quad G(0, x)(u_1, u_2) = (e^{2\pi i x} u_1, e^{-2\pi i x} u_2), \\ G(1, x)(u_1, u_2) = (u_1, u_2), \quad G(y, 0) = G(y, 1) \quad \forall x, y \in [0, 1].$$

Solution: We can take

$$(23.36) \quad G(y, x) = U(-y) \begin{pmatrix} \text{Id} & 0 \\ 0 & e^{-2\pi i x} \end{pmatrix} U(y) \begin{pmatrix} e^{2\pi i x} & 0 \\ 0 & \text{Id} \end{pmatrix}.$$

By the same reasoning as above, this is an homotopy of closed curves of invertible operators on  $H \oplus H$  which satisfies (23.35).

*Problem P10.6* Now, think about combining the various constructions above in the following way. Show that on  $l_2(H)$  there is an homotopy like (23.34),  $\tilde{G} : [0, 1]^2 \longrightarrow \text{GL}(l_2(H))$ , (very like in fact) such that

$$(23.37) \quad \tilde{G}(0, x) \{u_k\}_{k=1}^\infty = \{\exp((-1)^k 2\pi i x) u_k\}_{k=1}^\infty, \\ \tilde{G}(1, x) = \text{Id}, \quad \tilde{G}(y, 0) = \tilde{G}(y, 1) \quad \forall x, y \in [0, 1].$$

Solution: We can divide  $l_2(H)$  into its odd and even parts

$$(23.38) \quad D : l_2(H) \ni v \longmapsto (\{v_{2i-1}\}, \{v_{2i}\}) \in l_2(H) \oplus l_2(H) \longleftarrow H \oplus H.$$

and then each copy of  $l_2(H)$  on the right with  $H$  (using the same isometric isomorphism). Then the homotopy in the previous problem is such that

$$(23.39) \quad \tilde{G}(x, y) = D^{-1}G(y, x)D$$

accomplishes what we want.

*Problem P10.7: Eilenberg's swindle* For an infinite dimensional separable Hilbert space, construct an homotopy – meaning a continuous map  $G : [0, 1]^2 \longrightarrow \text{GL}(H)$  – with  $G(0, x) = L(x)$  in (23.28) and  $G(1, x) = \text{Id}$  and of course  $G(y, 0) = G(y, 1)$  for all  $x, y \in [0, 1]$ .

Hint: Just put things together – of course you can rescale the interval at the end to make it all happen over  $[0, 1]$ . First ‘divide  $H$  into 2 copies of itself’ and deform from  $L$  to  $M(1, x)$  in (23.30). Now, ‘divide the second  $H$  up into  $l_2(H)$ ’ and apply an argument just like the preceding problem to turn the identity on this factor into alternating terms multiplying by  $\exp(\pm 4\pi i x)$  – starting with  $-$ . Now, you are on  $H \oplus l_2(H)$ , ‘renumbering’ allows you to regard this as  $l_2(H)$  again and when you do so your curve has become alternate multiplication by  $\exp(\pm 4\pi i x)$  (with  $+$  first). Finally then, apply the preceding problem again, to deform to the identity (always of course through closed curves). Presto, Eilenberg's swindle!

Solution: By rescaling the variables above, we now have three homotopies, always through periodic families. On  $H \oplus H$  between  $L(x) = e^{2\pi i x} \text{Id}$  and the matrix

$$(23.40) \quad \begin{pmatrix} e^{4\pi i x} \text{Id} & 0 \\ 0 & \text{Id} \end{pmatrix}.$$

Then on  $H \oplus l_2(H)$  we can deform from

$$(23.41) \quad \begin{pmatrix} e^{4\pi i x} \text{Id} & 0 \\ 0 & \text{Id} \end{pmatrix} \text{ to } \begin{pmatrix} e^{4\pi i x} \text{Id} & 0 \\ 0 & \tilde{G}(0, x) \end{pmatrix}$$

with  $\tilde{G}(0, x)$  in (23.37). However we can then identify

$$(23.42) \quad H \oplus l_2(H) = l_2(H), \quad (u, v) \longmapsto w = \{w_j\}, \quad w_1 = u, \quad w_{j+1} = v_j, \quad j \geq 1.$$

This turns the matrix of operators in (23.41) into  $\tilde{G}(0, x)^{-1}$ . Now, we can apply the same construction to deform this curve to the identity. Notice that this really does ultimately give an homotopy, which we can renormalize to be on  $[0, 1]$  if you insist, of curves of operators on  $H$  – at each stage we transfer the homotopy back to  $H$ .