

Course 18.327 and 1.130 Wavelets and Filter Banks

**Modulation and Polyphase
Representations:
Noble Identities;
Block Toeplitz Matrices
and Block z-transforms;
Polyphase Examples**

Modulation Matrix

Matrix form of PR conditions:

$$[F_0(z) \ F_1(z)] \underbrace{\begin{bmatrix} H_0(z) & H_0(-z) \\ H_1(z) & H_1(-z) \end{bmatrix}}_{\text{Modulation matrix, } H_m(z)} = [2z^{-\ell} \ 0]$$

Modulation matrix, $H_m(z)$

So

$$[F_0(z) \ F_1(z)] = [2z^{-\ell} \ 0] H_m^{-1}(z)$$

$$H_m^{-1}(z) = \frac{1}{\Delta} \begin{bmatrix} H_1(-z) & -H_0(-z) \\ -H_1(z) & H_0(z) \end{bmatrix}$$

$$\Delta = H_0(z) H_1(-z) - H_0(-z) H_1(z) \quad (\text{must be non-zero})$$

$$\Rightarrow \left. \begin{aligned} F_0(z) &= \frac{1}{\Delta} 2z^{-\ell} H_1(-z) \\ F_1(z) &= \frac{1}{\Delta} 2z^{-\ell} H_0(-z) \end{aligned} \right\} \text{Require these to be FIR}$$

Suppose we choose $\Delta = 2z^{-\ell}$
Then

$$\left. \begin{aligned} F_0(z) &= H_1(-z) \\ F_1(z) &= -H_0(-z) \end{aligned} \right\}$$

3

Synthesis modulation matrix:

Complete the second row of matrix PR conditions by replacing z with $-z$:

$$\underbrace{\begin{bmatrix} F_0(z) & F_1(z) \\ F_0(-z) & F_1(-z) \end{bmatrix}}_{\text{Synthesis modulation matrix, } F_m(z)} \begin{bmatrix} H_0(z) & H_0(-z) \\ H_1(z) & H_1(-z) \end{bmatrix} = 2 \begin{bmatrix} z^{-\ell} & 0 \\ 0 & (-z)^{-\ell} \end{bmatrix}$$


Synthesis modulation matrix, $F_m(z)$

Note the transpose convention in $F_m(z)$.

4

Noble Identities

1. Consider




$$U(z) = H(z^2)X(z)$$

$$Y(z) = \frac{1}{2} \{U(z^{1/2}) + U(-z^{1/2})\} \quad (\text{downsampling})$$

$$= \frac{1}{2} \{H(z)X(z^{1/2}) + H(z)X(-z^{1/2})\}$$


$$= H(z) \cdot \frac{1}{2} \{X(z^{1/2}) + X(-z^{1/2})\} \Rightarrow \text{can downsample first}$$

First Noble identity:



5

2. Consider




$$U(z) = H(z)X(z)$$

$$Y(z) = U(z^2) \quad (\text{upsampling})$$

$$= H(z^2)X(z^2) \Rightarrow \text{can upsample first}$$


Second Noble Identity:



6


Derivation of Polyphase Form

1. Filtering and downsampling:





$$H(z) = H_{\text{even}}(z^2) + z^{-1} H_{\text{odd}}(z^2); \quad h_{\text{even}}[n] = h[2n]$$

$$h_{\text{odd}}[n] = h[2n+1]$$




7




8



2. Upsampling and filtering



$$F(z) = F_{\text{even}}(z^2) + z^{-1} F_{\text{odd}}(z^2)$$



9



**Polyphase
Form**

10

Polyphase Matrix

Consider the matrix corresponding to the analysis filter bank in interleaved form. This is a block Toeplitz matrix:

$$H_b = \begin{bmatrix} \vdots & & & & & & \\ \dots & \boxed{h_0[3] \ h_0[2]} & \boxed{h_0[1] \ h_0[0]} & 0 & 0 & \dots \\ \dots & \boxed{h_1[3] \ h_1[2]} & \boxed{h_1[1] \ h_1[0]} & 0 & 0 & \dots \\ \dots & 0 & 0 & \boxed{h_0[3] \ h_0[2]} & \boxed{h_0[1] \ h_0[0]} & \dots \\ \dots & 0 & 0 & \boxed{h_1[3] \ h_1[2]} & \boxed{h_1[1] \ h_1[0]} & \dots \\ & & & \vdots & & \end{bmatrix}$$

4-tap Example

11

Taking block z-transform we get:

$$\begin{aligned} H_p(z) &= \begin{bmatrix} h_0[0] & h_0[1] \\ h_1[0] & h_1[1] \end{bmatrix} + z^{-1} \begin{bmatrix} h_0[2] & h_0[3] \\ h_1[2] & h_1[3] \end{bmatrix} \\ &= \begin{bmatrix} h_0[0] + z^{-1} h_0[2] & h_0[1] + z^{-1} h_0[3] \\ h_1[0] + z^{-1} h_1[2] & h_1[1] + z^{-1} h_1[3] \end{bmatrix} \\ &= \begin{bmatrix} H_{0,\text{even}}(z) & H_{0,\text{odd}}(z) \\ H_{1,\text{even}}(z) & H_{1,\text{odd}}(z) \end{bmatrix} \end{aligned}$$

This is the polyphase matrix for a 2-channel filter bank.

12

Similarly, for the synthesis filter bank:

$$F_b = \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ \boxed{\begin{matrix} f_0[0] & f_1[0] \\ f_0[1] & f_1[1] \end{matrix}} & \begin{matrix} 0 & 0 \\ 0 & 0 \end{matrix} \\ \cdots & \boxed{\begin{matrix} f_0[2] & f_1[2] \\ f_0[3] & f_1[3] \end{matrix}} & \begin{matrix} f_0[0] & f_1[0] \\ f_0[1] & f_1[1] \end{matrix} & \cdots \\ \begin{matrix} 0 & 0 \\ 0 & 0 \end{matrix} & \begin{matrix} f_0[2] & f_1[2] \\ f_0[3] & f_1[3] \end{matrix} \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

13

$$F_p(z) = \begin{bmatrix} f_0[0] & f_1[0] \\ f_0[1] & f_1[1] \end{bmatrix} + z^{-1} \begin{bmatrix} f_0[2] & f_1[2] \\ f_0[3] & f_1[3] \end{bmatrix}$$

$$= \begin{bmatrix} F_{0,\text{even}}[z] & F_{1,\text{even}}[z] \\ F_{0,\text{odd}}[z] & F_{1,\text{odd}}[z] \end{bmatrix}$$

Note transpose convention for synthesis polyphase matrix

- Perfect reconstruction condition in polyphase domain:

$$F_p(z) H_p(z) = I \quad (\text{centered form})$$

This means that $H_p(z)$ must be invertible for all z on the unit circle, i.e.

$$\det H_p(e^{j\omega}) \neq 0 \text{ for all frequencies } \omega.$$

14

- Given that the analysis filters are FIR, the requirement for the synthesis filters to be also FIR is:

$$\det H_p(z) = z^{-\ell} \quad (\text{simple delay})$$

because $H_p^{-1}(z)$ must be a polynomial.

- Condition for orthogonality: $F_p(z)$ is the transpose of $H_p(z)$, i.e.

$$H_p^T(z^{-1}) H_p(z) = I$$

i.e. $H_p(z)$ should be paraunitary.

15

Relationship between Modulation and Polyphase Matrices

$$H_0(z) = H_{0,\text{even}}(z^2) + z^{-1} H_{0,\text{odd}}(z^2); \quad \begin{cases} h_{0,\text{even}}[n] = h_0[2n] \\ h_{0,\text{odd}}[n] = h_0[2n+1] \end{cases}$$

$$H_1(z) = H_{1,\text{even}}(z^2) + z^{-1} H_{1,\text{odd}}(z^2)$$

Two more equations by replacing z with $-z$.

So in matrix form:

$$\underbrace{\begin{bmatrix} H_0(z) & H_0(-z) \\ H_1(z) & H_1(-z) \end{bmatrix}}_{H_m(z)} = \underbrace{\begin{bmatrix} H_{0,\text{even}}(z^2) & H_{0,\text{odd}}(z^2) \\ H_{1,\text{even}}(z^2) & H_{1,\text{odd}}(z^2) \end{bmatrix}}_{H_p(z^2)} \begin{bmatrix} 1 & 1 \\ z^{-1} & -z^{-1} \end{bmatrix}$$

Modulation matrix Polyphase matrix

16

But

$$\begin{bmatrix} 1 & 1 \\ z^{-1} & -z^{-1} \end{bmatrix} = \underbrace{\begin{bmatrix} 1 & \\ & z^{-1} \end{bmatrix}}_{D_2(z)} \underbrace{\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}}_{F_2}$$

Delay Matrix 2-point DFT Matrix

$$F_N = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & w & w^2 & \dots & w^{N-1} \\ 1 & w^2 & w^4 & \dots & w^{2(N-1)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & w^{N-1} & w^{2(N-1)} & \dots & w^{(N-1)^2} \end{bmatrix}; \quad w = e^{\frac{2\pi}{N}} \rightarrow \text{N-point DFT Matrix}$$

$$F_N^{-1} = \frac{1}{N} \overline{F_N}$$

↑ Complex conjugate: replace w with $\overline{w} = e^{-\frac{2\pi}{N}}$

17

So, in general

$$H_m(z) F_N^{-1} = H_p(z^N) D_N(z)$$

N = # of channels in filterbank
($N = 2$ in our example)

18

Polyphase Matrix

Example: Daubechies 4-tap filter

$$h_0[0] = \frac{1+\sqrt{3}}{4\sqrt{2}} \quad h_0[1] = \frac{3+\sqrt{3}}{4\sqrt{2}} \quad h_0[2] = \frac{3-\sqrt{3}}{4\sqrt{2}} \quad h_0[3] = \frac{1-\sqrt{3}}{4\sqrt{2}}$$

$$H_0(z) = \frac{1}{4\sqrt{2}} \{(1 + \sqrt{3}) + (3 + \sqrt{3}) z^{-1} + (3 - \sqrt{3}) z^{-2} + (1 - \sqrt{3}) z^{-3}\}$$

$$H_1(z) = \frac{1}{4\sqrt{2}} \{(1 - \sqrt{3}) - (3 - \sqrt{3}) z^{-1} + (3 + \sqrt{3}) z^{-2} - (1 + \sqrt{3}) z^{-3}\}$$

19

Time domain:

$$\begin{aligned} h_0[0]^2 + h_0[1]^2 + h_0[2]^2 + h_0[3]^2 &= \frac{1}{32} \{(4 + 2\sqrt{3}) + (12 + 6\sqrt{3}) + \\ &\quad (12 - 6\sqrt{3}) + (4 - 2\sqrt{3})\} \\ &= 1 \end{aligned}$$

$$h_0[0] h_0[2] + h_0[1] h_0[3] = \frac{1}{32} \{(2\sqrt{3}) + (-2\sqrt{3})\}$$

$$= 0$$

i.e. filter is orthogonal to its double shifts

20

Polyphase Domain:

$$H_{0,\text{even}}(z) = \frac{1}{4\sqrt{2}} \{(1 + \sqrt{3}) + (3 - \sqrt{3}) z^{-1}\}$$

$$H_{0,\text{odd}}(z) = \frac{1}{4\sqrt{2}} \{(3 + \sqrt{3}) + (1 - \sqrt{3}) z^{-1}\}$$

$$H_{1,\text{even}}(z) = \frac{1}{4\sqrt{2}} \{(1 - \sqrt{3}) + (3 + \sqrt{3}) z^{-1}\}$$

$$H_{1,\text{odd}}(z) = \frac{1}{4\sqrt{2}} \{-(3 - \sqrt{3}) - (1 + \sqrt{3}) z^{-1}\}$$

$$H_p(z) = \frac{1}{4\sqrt{2}} \underbrace{\begin{bmatrix} 1 + \sqrt{3} & 3 + \sqrt{3} \\ 1 - \sqrt{3} & -(3 - \sqrt{3}) \end{bmatrix}}_A + \frac{1}{4\sqrt{2}} \underbrace{\begin{bmatrix} 3 - \sqrt{3} & 1 - \sqrt{3} \\ 3 + \sqrt{3} & -(1 + \sqrt{3}) \end{bmatrix}}_B z^{-1}$$

21

$$H_p(z) = A + B z^{-1}$$

$$\begin{aligned} H_p^T(z^{-1}) H_p(z) &= (A^T + B^T z)(A + Bz^{-1}) \\ &= (A^T A + B^T B) + A^T B z^{-1} + B^T A z \end{aligned}$$

$$\begin{aligned} A^T A &= \frac{1}{4\sqrt{2}} \begin{bmatrix} 1 + \sqrt{3} & 1 - \sqrt{3} \\ 3 + \sqrt{3} & -(3 - \sqrt{3}) \end{bmatrix} \frac{1}{4\sqrt{2}} \begin{bmatrix} 1 + \sqrt{3} & 3 + \sqrt{3} \\ 1 - \sqrt{3} & -(3 - \sqrt{3}) \end{bmatrix} \\ &= \frac{1}{32} \begin{bmatrix} (4 + 2\sqrt{3}) + (4 - 2\sqrt{3}) & (6 + 4\sqrt{3}) - (6 - 4\sqrt{3}) \\ (6 + 4\sqrt{3}) - (6 - 4\sqrt{3}) & (12 + 6\sqrt{3}) + (12 - 6\sqrt{3}) \end{bmatrix} \\ &= \begin{bmatrix} 1/4 & \sqrt{3}/4 \\ \sqrt{3}/4 & 3/4 \end{bmatrix} \end{aligned}$$

22

$$\begin{aligned}
\mathbf{B}^T \mathbf{B} &= \frac{1}{4\sqrt{2}} \begin{bmatrix} 3 - \sqrt{3} & 3 + \sqrt{3} \\ 1 - \sqrt{3} & -(1 + \sqrt{3}) \end{bmatrix} \frac{1}{4\sqrt{2}} \begin{bmatrix} 3 - \sqrt{3} & 1 - \sqrt{3} \\ 3 + \sqrt{3} & -(1 + \sqrt{3}) \end{bmatrix} \\
&= \frac{1}{32} \begin{bmatrix} (12 - 6\sqrt{3}) + (12 + 6\sqrt{3}) & (6 - 4\sqrt{3}) - (6 + 4\sqrt{3}) \\ (6 - 4\sqrt{3}) - (6 + 4\sqrt{3}) & (4 - 2\sqrt{3}) + (4 + 2\sqrt{3}) \end{bmatrix} \\
&= \begin{bmatrix} 3/4 & -\sqrt{3}/4 \\ -\sqrt{3}/4 & 1/4 \end{bmatrix}
\end{aligned}$$

$$\Rightarrow \mathbf{A}^T \mathbf{A} + \mathbf{B}^T \mathbf{B} = \mathbf{I}$$

23

$$\begin{aligned}
\mathbf{A}^T \mathbf{B} &= \frac{1}{4\sqrt{2}} \begin{bmatrix} 1 + \sqrt{3} & 1 - \sqrt{3} \\ 3 + \sqrt{3} & -(3 - \sqrt{3}) \end{bmatrix} \frac{1}{4\sqrt{2}} \begin{bmatrix} 3 - \sqrt{3} & 1 - \sqrt{3} \\ 3 + \sqrt{3} & -(1 + \sqrt{3}) \end{bmatrix} \\
&= \frac{1}{32} \begin{bmatrix} (2\sqrt{3}) + (-2\sqrt{3}) & (-2) - (-2) \\ (6) - (6) & (-2\sqrt{3}) + (2\sqrt{3}) \end{bmatrix} \\
&= \mathbf{0}
\end{aligned}$$

$$\mathbf{B}^T \mathbf{A} = (\mathbf{A}^T \mathbf{B})^T = \mathbf{0}$$

So

$$\mathbf{H}_p^T(z^{-1}) \mathbf{H}_p(z) = \mathbf{I} \quad \text{i.e. } \mathbf{H}_p(z) \text{ is a Paraunitary Matrix}$$

24

Modulation domain:

$$H_0(z) H_0(z^{-1}) = P(z) = \frac{1}{16} (-z^3 + 9z + 16 + 9z^{-1} - z^{-3})$$

$$H_0(-z) H_0(-z^{-1}) = P(-z) = \frac{1}{16} (z^3 - 9z + 16 - 9z^{-1} + z^{-3})$$

So


$$H_0(z) H_0(z^{-1}) + H_0(-z) H_0(-z^{-1}) = 2$$

i.e.

$$|H_0(\omega)|^2 + |H_0(\omega + \pi)|^2 = 2$$


25

Magnitude Response of Daubechies 4-tap filter.



26

Phase response of Daubechies 4-tap filter.



27